arXiv:1108.5407v1 [cond-mat.other] 27 Aug 2011

Superfluidity in Two-Dimensional Imbalanced Fermi Gases

Heron Caldas,[1, *] A. L. Mota,[1] R. L. S. Farias,[1] and L. A. Souza[1]
[1] *Departamento de Ciências Naturais, Universidade Federal de São João del Rei,*
36301-160, São João del Rei, MG, Brazil

We study the zero temperature ground state of a two-dimensional atomic Fermi gas with chemical potential and population imbalance in the mean-field approximation. All calculations are performed in terms of the two-body binding energy ϵ_B, whose variation allows to investigate the evolution from the BEC to the BCS regimes. By means of analytical and exact expressions we show that, similarly to what is found in three dimensions, at fixed chemical potentials, BCS is the ground state until the critical imbalance h_c after which there is a first-order phase transition to the normal state. We also show that for a fixed ratio ϵ_B/ϵ_F, where ϵ_F is the two-dimensional Fermi energy, as the density imbalance m is increased from zero, the ground state evolves from BCS to phase separation to the normal state. At the critical imbalance m_c phase separation is not supported and the normal phase is energetically preferable. The BCS-BEC crossover is discussed in balanced and imbalanced configurations. Possible pictures of what may be found experimentally in these systems are also shown.

PACS numbers: 71.30.+h,36.20.Kd,11.10.Kk

I. INTRODUCTION

In the last few years, great experimental advances have permitted the manipulation of trapped neutral ultracold two-spin-components atomic Fermi gases with tremendous accuracy. The essential technics under full domain are the cooling, trapping, the control of the number of atoms in the sample, and the tunning of the interatomic (s-wave) interactions via the application of an external uniform magnetic (Feshbach resonance) field [1]. This allowed the investigation of the crossover from the Bardeen-Cooper-Schrieffer (BCS) phase of weakly bound Cooper pairs to the Bose-Einstein condensate (BEC) phase of strongly bound diatomic molecules in three-dimensional (3D) trapped Fermi gases [2–4].

These many-body quantum gases are even more exciting when the two-components have mass, chemical potential or population imbalance, since this situation can be met in many areas of physics, from condensed matter to high-density quark matter. Many exotic phases have been proposed as the ground state of these imbalanced Fermi gases, such as the homogeneous Sarma [5] or breached-pair state [6], the Fulde-Ferrel-Larkin-Ovchinnikov (FFLO) state with modulated order parameter [7], phase separation of superfluid and normal components in real space [8, 9], deformed Fermi surfaces [10], and magnetized superfluid in the BEC side of the resonance [11]. Recent experiments in imbalanced atomic Fermi gases confined in 3D harmonic traps have observed that these systems phase separate into a unpolarized superfluid core surrounded by a normal outer region [12–16].

Another very important parameter under control is the trap geometry, which permits the experimental investigation of fermionic atoms in lower dimensions. A two-dimensional (2D) trapped Fermi gas may be achieved in experiments by flattening magnetic or dipolar confinements [17], by trapping atoms in specially designed pancake potentials [18], by radio frequency (rf)-induced two-dimensional traps [19], or by gravito-optical surface traps [20].

Motivated by these experimental progresses, some recent papers have investigated two-component Fermi gases in 2D [21–24]. One-dimentional (1D) two-component Fermi systems have also received attention recently both theoretically [25, 26] and experimentally [27]. Besides, spin imbalance has also been investigated in the context of metallic systems as, for example, electron-hole bilayer [28, 29] and multi-band [30] systems.

In this paper we study the zero temperature (T) ground state of a 2D atomic Fermi gas with chemical potential and population imbalance in the mean-field approximation. We show that in the grand canonical ensemble the fundamental state is BCS up to a critical value of the chemical potentials asymmetry h_c, after which there is a quantum phase transition to the normal state. In the canonical ensemble we study in detail the issue of phase separation. We show that the normal phase is always unstable to phase separation when the density asymmetry m is less than the critical imbalance m_c. All calculations are performed in terms of the two-body binding energy ϵ_B, whose variation allows to investigate the evolution from the BEC to the BCS regimes. Since the recent 3D trap experiments of Shin et al. [17] have not observed the FFLO state, we do not consider in this work the possibility of this phase in 2D. The observation of the FFLO state is indeed an experimental challenging, since it is predicted to exist only in a narrow window of asymmetry between the particle's chemical potentials. Nevertheless it is worth to mention that significant FFLO-like correlations are expected to be present in 1D systems for a range of polarizations [27].

---

[*] Electronic address: hcaldas@ufsj.edu.br

The main results of this paper may be summarized as:
I. We show that in a zero T imbalanced 2D gas of fermionic atoms, BCS is the ground state up to chemical potentials imbalance h_c at which there is a first-order phase transition to the normal state.
II. We demonstrate that, as found theoretically and experimentally in 3D, for a fixed two-body binding energy e_B, as the imbalance m is increased from zero the stable states are BCS, phase separation and normal phase.
III. For balanced systems, the BCS-BEC crossover is governed only by e_B. However, for imbalanced systems, the BCS-BEC crossover will depend on e_B and the imbalance m. For a fix value of e_B, the crossover is possible only for asymmetries supported by phase separation. Above a critical value of the number densities imbalance m_c, phase separation is destroyed (in favor of the normal phase) as well as the path to BEC regime.

The paper is organized as follows: In Sec. II we present the model Hamiltonian and derive an analytical expression for the zero temperature thermodynamical potential in the mean-field approximation for the two-component gas of fermionic atoms in 2D. In Sec. III we employ this expression to investigate the stability of the possible phases of 2D imbalanced Fermi gases, both at fixed chemical potentials and densities. We also discuss the BCS-BEC crossover in balanced and imbalanced systems. Sec. IV is devoted to the conclusions.

II. THE MODEL HAMILTONIAN AND THE MEAN-FIELD THEORY

We start by considering a 2D nonrelativistic dilute system of fermionic atoms of mass M, with two hyperfine states labeled as $\sigma =\uparrow, \downarrow$. This spin $\uparrow$ and $\downarrow$ mixture could, in principle, be done with the two lowest hyperfine states of ^{6}Li atoms, as in the 3D experiments [13, 14, 16]. Their single-particle dispersion relations are given by $\xi_k = \frac{\hbar^2 k^2}{2M}$. Throughout the paper we set $\hbar = 1$. Since at low densities (appropriate to describe the ultracold trapped Fermi gases we are interested) the form of the potential is not probed, it can be considered that the atoms interact via a contact interaction which is modeled by the following pairing Hamiltonian:

$$H = H_0 + H_{\rm int}, \tag{1}$$

where

$$H_0 = \sum_{k,\sigma=\uparrow,\downarrow} \epsilon_k^\sigma \psi_\sigma^\dagger(k)\psi_\sigma(k), \tag{2}$$

is the kinetic (free) part of H and $H_{\rm int}$ is given below. $\psi_\sigma^\dagger(k)$ and $\psi_\sigma(k)$ in Eq. (2) are the creation and annihilation operators, respectively, for the $\uparrow, \downarrow$ particles. To assure population imbalance in the system, we have introduced different chemical potential for the species σ as $\mu_\sigma = \mu + \sigma h$, where $\sigma h \equiv \pm h$. Then, the chemical potential μ_σ fix the number densities n_σ of the different fermions. The new dispersions for the free species σ, relative to their Fermi energies, are $\epsilon_k^\sigma \equiv \xi_k - \mu_\sigma$. The interaction Hamiltonian is given by

$$H_{int} = g \sum_{k,k'} \psi_\uparrow^\dagger(k)\psi_\downarrow^\dagger(-k)\psi_\downarrow(-k')\psi_\uparrow(k'), \tag{3}$$

where the bare coupling constant g is negative, to express the attractive (s-wave) interaction between the spin $\uparrow$ and $\downarrow$ fermionic atoms.

After the mean field (MF) approximation and the subsequent diagonalization of the expression for H_{MF}, we arrive at the following expression for thermodynamic potential:

$$\begin{aligned} \Omega &= \int_{k<k_1,\ k>k_2} \frac{d^2k}{(2\pi)^2}(\epsilon_k^b - \varepsilon_k^\beta) - \frac{\Delta^2}{g} \\ &+ \int_{k_1}^{k_2} \frac{d^2k}{(2\pi)^2}\epsilon_k^b \\ &= \int_{k<k_1,\ k>k_2} \frac{d^2k}{(2\pi)^2}(\epsilon_k^+ - E_k) - \frac{\Delta^2}{g} \\ &+ \int_{k_1}^{k_2} \frac{d^2k}{(2\pi)^2}\epsilon_k^b, \end{aligned} \tag{4}$$

where, for simplicity of notation we have labeled $\downarrow = a$, $\uparrow = b$, and we have defined $\epsilon_k^\pm = \frac{\epsilon_k^a \pm \epsilon_k^b}{2}$, $\varepsilon_k^\beta = -\epsilon_k^- + E_k$, $E_k = \sqrt{{\epsilon_k^+}^2 + \Delta^2}$, and the constant pairing gap is given by $\Delta = -g\int \frac{d^2k}{(2\pi)^2}\langle \psi_\downarrow^\dagger(-k)\psi_\uparrow^\dagger(k)\rangle = \Delta^*$. In Eq. (4)

$$k_{1,2}^2 = \frac{k_a^2 + k_b^2}{2} \pm \frac{1}{2}\sqrt{(k_b^2 - k_a^2)^2 - 16M^2\Delta^2} \tag{5}$$

are the roots of ε_k^β, and $k_\alpha = \sqrt{2M\mu_\alpha}$ is the Fermi momentum of the species $\alpha = a, b$. From the equation above one sees that the condition for Δ such that $k_{1,2}$ is real is $\Delta \leq \frac{k_b^2 - k_a^2}{4M}$.

To regulate the ultraviolet divergence associated with the first integral in Eq. (4) we introduce [31]:

$$\frac{1}{U} = \int \frac{d^2k}{(2\pi)^2}\frac{1}{2\xi_k + |\epsilon_B|}, \tag{6}$$

where $U \equiv -g > 0$, and ϵ_B is the 2D two-body binding energy. In order to make contact with current experiments, it is convenient to relate ϵ_B to the three dimensional scattering length a_s. In the scattering of atoms confined in the axial direction by a harmonic potential with characteristic frequency ω_L they are related by [21, 32]:

$$|\epsilon_B| = \frac{C\omega_L}{\pi} exp\left(\sqrt{2\pi}\frac{l_L}{a_s}\right), \tag{7}$$

where a_s is the 3D s-wave scattering length, $\omega_L = \sqrt{8\pi^2 V_0/(M\lambda^2)}$, $l_L = 1/\sqrt{M\omega_L}$, and $C \approx 0.915$. V_0 is the amplitude of the periodic potential $V_0 \sin^2(2\pi z/\lambda)$ generated by two counter-propagating laser beans with length λ parallel to the z-axis [21].

Thus we obtain an analytical expression for the grand canonical thermodynamic potential at $T = 0$:

$$\begin{aligned}\bar{\Omega} &= \Omega(\Delta,\mu) + \Theta_h \Omega(\Delta,h) \qquad (8)\\ &= \Delta^2\left[\ln\left(\frac{\sqrt{\mu^2+\Delta^2}-\mu}{|\epsilon_B|}\right) - \frac{1}{2}\right] - \mu\left(\sqrt{\mu^2+\Delta^2}+\mu\right) - \Theta_h\left[2h\sqrt{h^2-\Delta^2} - \Delta^2\ln\left(\frac{h+\sqrt{h^2-\Delta^2}}{h-\sqrt{h^2-\Delta^2}}\right)\right],\end{aligned}$$

where $\bar{\Omega} \equiv \frac{4\pi}{M}\Omega$, $\Theta_h \equiv \Theta(h^2 - \Delta^2)$, $\mu = \frac{\mu_a+\mu_b}{2}$, $h = \frac{\mu_b-\mu_a}{2}$, and $\Theta(x)$ is the Heaviside step function, defined as 1 if $x \geq 0$, and 0 if $x < 0$. With these definitions, Eq. (5) may be written as $k_{1,2}^2 = 2M(\mu \pm \sqrt{h^2-\Delta^2})$ and the condition for real $k_{1,2}^2$ is now $\Delta \leq h$.

III. PHASES OF 2D IMBALANCED FERMI SYSTEMS AND THEIR STABILITIES

A. Fixed Chemical Potentials

In the grand canonical ensemble the chemical potentials of the two-components are held fixed. This happens when the system is connected to reservoirs of species a and b such that the particle densities are allowed to change. We analyze the ground state of this case for balanced and imbalanced configurations.

1. *Balanced Systems*

The gap and number equations are obtained by $\partial\Omega/\partial\Delta = 0$ and $n_\alpha = -\partial\Omega/\partial\mu_\alpha$, respectively. For the balanced system where $h = 0$, which implies $\Theta_h = 0$, we find

$$\sqrt{\mu^2+\Delta^2} - \mu = |\epsilon_B|, \qquad (9)$$

and

$$\sqrt{\mu^2+\Delta^2} + \mu = 2\epsilon_F, \qquad (10)$$

where the two dimensional Fermi energy is defined as $\epsilon_F = \frac{\pi n_T}{M}$, with $n_T = n_a + n_b$, see Appendix A. In the balanced configuration $n_a = n_b \equiv n = \frac{n_T}{2}$. Solving these two equations self-consistently we arrive at

$$\Delta_0 = \sqrt{2\epsilon_F|\epsilon_B|}, \qquad (11)$$

and

$$\mu_0 = \epsilon_F - \frac{|\epsilon_B|}{2}. \qquad (12)$$

The value of the free energy at the minimum is

$$\Omega(h=0, \Delta=\Delta_0) \equiv \Omega_0 = -\kappa\left(\mu_0 + \frac{|\epsilon_B|}{2}\right)^2, \qquad (13)$$

where $\kappa \equiv \frac{M}{2\pi}$, whereas the energy of the balanced normal state is given by

$$\Omega(h=\Delta=0) \equiv \Omega_b^N = -\kappa\mu_0^2. \qquad (14)$$

A direct comparison between Eqs. (13) and (14) shows that the superfluid state is energetically preferable to the normal state for any $\epsilon_B \neq 0$. Since a two-body bound state exists even for an arbitrarily small attraction in 2D [31], the pairing instability will always happens in 2D balanced two-component Fermi systems at $T = 0$.

2. *BCS-BEC Crossover in Balanced Systems*

The two well known [31, 33] equations (11) and (12) reveal that the BCS-BEC crossover can be accessed by varying ϵ_B from weak to strong interaction regimes. The BCS state is reached in the weak attraction limit or high density (ϵ_F), $|\epsilon_B| << \epsilon_F$, where $\mu_0 \cong \epsilon_F$ and the energy gap is found to be

$$\Delta_0 = 2\sqrt{\epsilon_F \mathrm{E}_\Lambda}\, e^{-\frac{2\pi}{MU}} << \epsilon_F, \qquad (15)$$

where $\mathrm{E}_\Lambda \equiv \frac{\Lambda^2}{2\mathrm{M}}$, and Λ is a momentum cutoff. To obtain Eq. (15) from Eq. (11) we have made use of Eq. (6). The opposite limit, of very strong attraction or low density, $|\epsilon_B| >> \epsilon_F$, results in the formation of composite bosons with $\mu_0 \cong -|\epsilon_B|/2$. Since the BCS and BEC regions are characterized by positive and negative chemical potentials, respectively, a clear distinction between these two regimes is the value of the binding energy at which μ_0 changes sign i.e., $|\epsilon_B| = 2\epsilon_F$.

3. Imbalanced Systems

We now turn our attention to the cases where $h \neq 0$. The free energy of the imbalanced normal state, $\Omega(h, \Delta = 0) \equiv \Omega_i^N$, is found to be

$$\Omega_i^N = \int_{k>k_2,k<k_1} \frac{d^2k}{(2\pi)^2}(\epsilon_k^+ - |\epsilon_k^+|) + \int_{k_1}^{k_2} \frac{d^2k}{(2\pi)^2}\epsilon_k^b. \tag{16}$$

In the limit $\Delta \to 0$ we find $k_1(\Delta = 0) = \sqrt{2M(\mu - h)} = k_F^a < k_F$ and $k_2(\Delta = 0) = \sqrt{2M(\mu + h)} = k_F^b > k_F$, where $k_F = \sqrt{2M\mu}$. We are considering μ_α and h positive. Thus the equation above is written as:

$$\Omega_i^N = \int_0^{k_F^a} \frac{d^2k}{(2\pi)^2}\epsilon_k^a + \int_0^{k_F^b} \frac{d^2k}{(2\pi)^2}\epsilon_k^b, \tag{17}$$

which gives, after the integration in k, the free energy of a (normal) two species gas of fermionic atoms in two dimensions

$$\begin{aligned} \Omega_i^N(\mu_a, \mu_b) &= -\frac{\kappa}{2}[(\mu - h)^2 + (\mu + h)^2] \\ &= -\frac{\kappa}{2}[\mu_a^2 + \mu_b^2]. \end{aligned} \tag{18}$$

The gap and number equation now read

$$\ln\left(\frac{\sqrt{\mu^2 + \Delta^2} - \mu}{|\epsilon_B|}\right) + \Theta_h \ln\left(\frac{h + \sqrt{h^2 - \Delta^2}}{h - \sqrt{h^2 - \Delta^2}}\right) = 0, \tag{19}$$

$$\sqrt{\mu^2 + \Delta^2} + \mu = 2\epsilon_F. \tag{20}$$

Note that for $h < \Delta$ Eq. (19) is reduced to Eq. (9). Seeking now solutions where $h > \Delta$ we then have to solve

$$\frac{\sqrt{\mu^2 + \Delta^2} - \mu}{|\epsilon_B|} = \frac{h - \sqrt{h^2 - \Delta^2}}{h + \sqrt{h^2 - \Delta^2}}. \tag{21}$$

Solving these equations self-consistently we find

$$\Delta_S(h) = \sqrt{\Delta_0(2h - \Delta_0)}. \tag{22}$$

From the equation above we see that the "Sarma state" (as explained below) gap and the chemical potential imbalance exist in the ranges

$$0 \leq \Delta_S(h) \leq \Delta_0, \tag{23}$$

$$\frac{\Delta_0}{2} \leq h \leq \Delta_0, \tag{24}$$

where the upper limit is imposed by the existence of real $k_{1,2}^2$.

From the graphical analysis of Ω as a function of Δ for various asymmetries, shown in Fig. 1, one sees that the first curve, from top to bottom, is the usual balanced system with its minimum at Δ_0. Increasing the imbalance h and keeping μ positive (to guarantee that we are in the BCS regime), the minimum is still located at Δ_0 up to a maximum or critical imbalance h_c, after which there is a quantum phase transition to the normal state with $\Delta = 0$. h_c is find through the equality $\Omega_0 = \Omega_i^N$, which yields:

$$h_c^2 = \mu_0|\epsilon_B| + \left(\frac{\epsilon_B}{2}\right)^2, \tag{25}$$

from which one easily finds plugging in the equation above μ_0 from Eq. (12) in the BCS limit ($|\epsilon_B| << e_F$): $h_c = \frac{\Delta_0}{\sqrt{2}}$.

We have seen above that, as happens in 3D [8, 9], the BCS phase turns from stable, while $h < h_c$, to metastable, when $h > h_c$, in which case the normal phase is stable. Besides these two phases, there is an unstable phase, known as Sarma state, corresponding to a local maximum of Ω versus Δ, that is located between the BCS minimum at Δ_0 and the normal phase with $\Delta = 0$.



FIG. 1: The thermodynamic potential Ω as a function of Δ for various chemical potential asymmetries. The top curve is for $h = 0$. The imbalance is being increased from top to bottom and in all curves for $h < h_c$ the minimum is at Δ_0. The curve for $h = h_c$ has two minima. After h_c, at which there is a quantum phase transition for the normal state, $\Omega(\Delta = 0, h > h_c) < \Omega(\Delta_0, h \leq h_c)$.

B. Fixed Number Densities

In the previous subsection we discussed the stability of several phases with fixed chemical potentials μ_a and μ_b. We investigate now the situations where the number densities n_a and n_b are fixed, since it is appropriate to describe trapped gases in current experiments. It is convenient to introduce now the "magnetization" defined (see Appendix A) as $m = -\frac{d\Omega}{dh} = n_b - n_a$, which will be needed to describe the energy of the Sarma phase in terms of the number densities, as discussed below:

$$m = \frac{M}{\pi}\sqrt{h^2 - \Delta^2}. \tag{26}$$

Solving equations (20), (21) and (26) self-consistently, we find the gap $\Delta_S(m)$ and the "average" chemical potential μ^S of the Sarma phase

$$\Delta_S(m) = \sqrt{\Delta_0^2 - \frac{2\pi\Delta_0}{M}m}, \tag{27}$$

where Δ_0, the gap parameter of the balanced system, is given by Eq. (11), and

$$\mu_S = \epsilon_F - \frac{{\Delta_S}^2}{4\epsilon_F}. \tag{28}$$

Note that the equations above reduce to Eqs. (11) and (12), respectively, since in the limit $m \to 0$, $\Delta_S \to \Delta_0$, as it should be. From Eq. (27) it is easy to obtain the windows for the Sarma gap and m:

$$0 \le \Delta_S(m) \le \Delta_0, \tag{29}$$

$$0 \le m \le m_{max}, \tag{30}$$

where $m_{max} = \frac{M\Delta_0}{2\pi}$ is the maximum value for the density imbalance in the Sarma phase, which is easily obtained from Eq. (27).

The expression for the energy E (not the free energy or thermodynamic potential) of the homogeneous Sarma phase is obtained as:

$$E(n_T, m, \Delta) = \Omega(\mu, h, \Delta) + \mu_a n_a + \mu_b n_b. \tag{31}$$

To write $E = E(n_T, m)$, we express $\mu_a n_a + \mu_b n_b = \mu n_T + hm$, with $\mu = \mu(n_T)$, $h = h(m)$ to find

$$\begin{aligned} E(n_T, m, \Delta) &\equiv E^S(n_a, n_b) = \Omega(n_T, m, \Delta) + \mu n_T + hm \\ &= \frac{M}{4\pi}\left\{\Delta_S^2\left[\ln\left(\frac{\Delta_S^2}{\Delta_0^2}\right) - \frac{1}{2}\right] - 2\epsilon_F\left(\epsilon_F - \frac{\Delta_S^2}{4\epsilon_F}\right) - 2\sqrt{\bar{m}^2 + \Delta_S^2}\,\bar{m} + \Delta_S^2 \ln\left(\frac{\sqrt{\bar{m}^2 + \Delta_S^2} + \bar{m}}{\sqrt{\bar{m}^2 + \Delta_S^2} - \bar{m}}\right)\right\} \\ &+ \left(\epsilon_F - \frac{\Delta_S^2}{4\epsilon_F}\right) n_T + \sqrt{\bar{m}^2 + \Delta_S^2}\, m, \end{aligned} \tag{32}$$

where $\bar{m} \equiv \frac{\pi m}{M}$.

To investigate which one is the ground state of an imbalanced system with different densities n_a and n_b, we have to see which energy of the following states we are considering is smaller, the normal $E^N(n_a, n_b)$, see Eq. (36) below, the homogeneous Sarma $E^S(n_a, n_b)$, or the phase separation (PS) $E^{PS}(n_a, n_b)$ state. The PS is an inhomogeneous phase where given n_a and n_b densities in a trap, a fraction $1 - x$ of the 2D (real) space is in the BCS phase where both species have a common density $n_a^{BCS} = n_b^{BCS} = n$ and the rest of particles are in the normal phase occupying the fraction x around[1] the BCS core, with densities $\tilde{n}_a$ and $\tilde{n}_b$. The most favored composition minimizes:

$$E^{PS} = Min_{x,\tilde{n}}[(1 - x)E(n) + xE^N(\tilde{n}_a, \tilde{n}_b)], \tag{33}$$

where the space fraction is obviously limited to the range $0 \le x \le 1$. The energy of the balanced superfluid phase is found as

---

[1] A surface energy is related to the interface between the normal and superfluid phases [34–37]. The surface tension associated with this interface has been interpreted as responsible for deformations in highly elongated small samples [14, 15]. These results are not consistent with the observations reported in Ref. [16]. We neglect the surface energy in the present analysis. We intent to come to this issue again soon taking into account corrections beyond the mean-field approximation.

$$\begin{aligned} E(n) &= \Omega(\Delta, h=0, \mu) + \mu n_T \\ &= \frac{M}{4\pi}\left\{\Delta^2\left[\ln\left(\frac{\Delta^2}{\Delta_0^2}\right) - 1\right] + 2\epsilon_F^2\right\} \\ &= \frac{M\Delta^2}{4\pi}\left[\ln\left(\frac{\Delta^2}{\Delta_0^2}\right) - 1\right] + \frac{\pi(2n)^2}{2M}. \end{aligned} \tag{34}$$

Since $E(n)$ which enters E^{PS} has to be written at its minimum, the equation above turns out to be:

$$\begin{aligned} E(n) &= \frac{\pi {n_T}^2}{2M} - \frac{M\Delta_0^2(n)}{4\pi} \\ &= \frac{2\pi}{M}n^2 - |\epsilon_B|n, \end{aligned} \tag{35}$$

where the gap parameter of the BCS phase is given by Eq. (11) and written as $\Delta_0(n) = 2\sqrt{\frac{\pi}{M}|\epsilon_B|n}$.

The energy of the normal state entering in Eq. (33) is easy to find from Eq. (18) as

$$\begin{aligned} E^N(n_a, n_b) &= \Omega^N(n_a, n_b) + \mu_a n_b + \mu_a n_b \\ &= \frac{1}{2\kappa}(n_a^2 + n_b^2). \end{aligned} \tag{36}$$

Note that since $n_a = \frac{n_T - m}{2}$ and $n_b = \frac{n_T + m}{2}$, the equation above for $m = 0$ gives $E^N = \frac{\pi n_T^2}{2M}$ agreeing with the result of Eq. (34) in the limit $\Delta \to 0$.

As we mentioned before, the n_a and n_b particle densities are accommodated in the trap as

$$\begin{aligned} n_a &= x\tilde{n}_a + (1-x)\tilde{n}, \\ n_b &= x\tilde{n}_b + (1-x)\tilde{n}. \end{aligned} \tag{37}$$

Then we rewrite Eq. (36) as

$$E^N(\tilde{n}_a, \tilde{n}_b) = \frac{1}{2\kappa}\left[\left(\frac{n_a - (1-x)\tilde{n}}{x}\right)^2 + \left(\frac{n_b - (1-x)\tilde{n}}{x}\right)^2\right] \tag{38}$$

It is interesting to note that Eqs. (35) and (36) are easily obtained from Eq. (32) in the limits $m \to 0$ and $m \to m_{max}$, respectively.

In the minimization of E^{PS} we shall find $x \approx 0$ and $\tilde{n} \approx n_a$ (or n_b) if $n_b \approx n_a$, which means that the entire system is in the superfluid phase. In the other limit, if $n_b >> n_a$, the system will be in the normal state, so we shall find in the minimization of E^{PS} $x \approx 1$ and $\tilde{n} \approx 0$. In the intermediate cases the system will phase separate, i.e. if the imbalance is less than the critical value, as experiments have shown for 3D gases.

1. Phase Competition

There are some limiting cases where the comparison between the phase separation and Sarma states can be done analytically and exactly. The first one is obtained when $m = 0$, giving $E^S(n) = E^{PS}(n) = E^{BCS}(n)$. The second case happens when $m = m_{max}$, which reduces the Sarma phase to the normal phase with $\Delta = 0$. This is easily seen from Eqs. (30) and (29). Then we define $\Delta E = E^{PS} - E^N$, which is written as

$$\begin{aligned} \Delta E &= \left[\frac{\tilde{n}^2}{\kappa} - E(\tilde{n})\right](x-1) \\ &+ \frac{1}{2\kappa}\left[(n_T - \tilde{n})^2 - 2n_a n_b + \tilde{n}^2\right]\left(\frac{1}{x} - 1\right). \end{aligned} \tag{39}$$

Minimization with respect to x yields:

$$x_{min} = \sqrt{\frac{(n_T - \tilde{n})^2 - 2n_a n_b + \tilde{n}^2}{2(\tilde{n}^2 - \kappa E(\tilde{n}))}}. \tag{40}$$

An upper bound on ΔE can be obtained by setting the density of the BCS component of the PS state $\tilde{n} = n_a$ and writing $n_b = n_a + m$, where for low imbalance, $m < n_a, n_b$, Eq. (40) gives:

$$x_{min} = \frac{m}{\sqrt{\frac{M^2\Delta_0^2(n_a)}{4\pi^2} + \frac{3}{2}n_a^2}}, \tag{41}$$

and

$$\begin{aligned} \Delta E(x_{min}) &= -\left(\frac{M\Delta_0^2(n_a)}{4\pi} + \frac{3\pi}{2M}n_a^2\right) \\ &\times\ (1 - x_{min})^2 < 0. \end{aligned} \tag{42}$$

Unlike the three-dimensional case where the result obtained is an approximation, valid only for small asymmetries $m << n_a, n_b$, where terms of order $m^3/na^{4/3}$ have been neglected [8, 9], the result above is exact and valid for any $m < n_a, n_b$. These results agree with the numerical analysis made to construct the phase diagram depicted in Fig. (2).

Let us now show the results of the minimization of Eq (33) with respect to x and n for a given n_a and n_b. The results are shown in Fig. (2).

We show in Fig. (3) an illustration of what may be found experimentally in an imbalanced $2D$ gas of femionic atoms. The first picture represents the balanced system with $m = 0$. The following pictures represent a set of experiments with fixed $e_B/e_F = 0.0318$ with increasing imbalance. From the second picture to the fifth the system phase separates and the sixth picture is for the critical imbalance m_c representing the normal phase.



FIG. 2: The phase diagram of a zero temperature imbalanced gas of fermionic atoms in 2D. The vertical axis $|e_B|/e_F$ with $m/n_T = 0$, is the (balanced) BCS phase, whereas the horizontal line $|e_B|/e_F = 0$ represents the (imbalanced) normal phase. Note that for each value of $|e_B|/e_F$ phase separation is the ground state until m_c. Above this value phase separation is not energetically favorable, and the normal phase has lower energy.

2. BCS-BEC Crossover in Imbalanced Systems

Now we investigate the BCS-BEC crossover in an imbalanced gas of fermionic atoms. As we have seen, there are three possible stable phases with fixed number densities. For zero imbalance between the densities, the only phase present is BCS (for any value of the interaction between the spin up and down fermionic atoms) and the crossover is that discussed in Subsection III A 2, which is governed only by ϵ_B. For imbalances where phase separation is possible, the BCS-BEC crossover can be realized in the superfluid core of the separated phases, again varying ϵ_B in $\mu_0 = \frac{2\pi\tilde{n}}{M} - \frac{|\epsilon_B|}{2}$, where $\tilde{n}$ is the BCS density that minimizes the PS formation, as described above Eq. (33). For large enough asymmetries, phase separation is not supported and the entire system is in the normal phase with no possibility of pairing formation and the crossover to the BEC regime. Then, when $m \neq 0$ the BCS-BEC crossover can not be accessed only by varying the biding energy since the imbalance plays an important role now.

IV. CONCLUSIONS

We have investigated superfluidity in two-dimensional imbalanced Fermi gases. With exact expressions, we have constructed the zero temperature mean-field phase diagram of these systems. We show that in the grand canonical ensemble, BCS is the ground state up to a critical chemical potential imbalance h_c at which there is a first-order phase transition to the normal state. In the canonical ensemble, relevant to current experiments, we also show that, as found theoretically and experimentally in 3D, for a fixed two-body binding energy e_B, as the imbalance m is increased from zero the stable states are BCS, phase separation and normal phase. For balanced systems, the BCS-BEC crossover is governed only by e_B. However, for imbalanced systems, the BCS-BEC crossover will depend on e_B and the imbalance m. For a fix value of e_B, the crossover is possible only for asymmetries supported by phase separation. Above a critical value of the number densities imbalance m_c, phase separation is destroyed in favor of the normal phase, as well as the path to BEC regime. Thus, for a given $m < m_c$ such that the system phase separates into BCS and normal phases, the BEC regime can be accessed by varying e_B from the weak to strong couplings. A smaller and smaller circular core shall be visibly apparent, as e_B



FIG. 3: Evolution of the phases of an imbalanced gas with fixed $|e_B|/e_F = 0.0318$ as a function of increasing m/n_T. The first picture, with $m/n_T = 0$ represent the BCS phase. Increasing m/n_T the system phase separates (pictures 2 to 5), and increasing the imbalance even more, the system finally goes to the normal phase at the critical imbalance $m_c/n_T = 0.179$, as seen in the last picture.

is increased, signalizing the condensation of the Cooper pairs. This is a direct prediction of our results that could be tested experimentally. Another interesting phenomena that could also be verified experimentally in a 2D trapped imbalanced Fermi gas [38], emerges in the opposite range, that of very large imbalance $m > m_c$ or, in the unities of Fig. 2, $m/n_T \approx 1$. In this situation, the minority species a correspond to impurities imersed in the majority species b Fermi liquid. As the interaction $|e_B|/e_F$ is increased, the impurities a surround themselves with a localized cloud of environment b atoms, forming Fermi polarons [39].

V. ACKNOWLEDGMENTS

We thank Drs. M. W. Zwierlein and L. H. C. M. Nunes for enlightening discussions. We also thank Dr. R. G. Hulet for a critical reading of the manuscript and for valuable suggestions. H. C. and A. L. M. are partially supported by CNPq. The authors also acknowledge partial support from FAPEMIG.

VI. APPENDIX A: DERIVATION OF THE MAGNETIZATION AND TOTAL NUMBER EQUATIONS

The total derivative of $\Omega = \Omega(\mu_a, \mu_b, \Delta)$ is written as

$$d\Omega = \frac{\partial\Omega}{\partial\mu_a}d\mu_a + \frac{\partial\Omega}{\partial\mu_b}d\mu_b + \frac{\partial\Omega}{\partial\Delta}d\Delta. \tag{43}$$

Since the densities have to evaluated at the minimum of Ω, i.e., at Δ_0, the last term vanishes. The particle densities are given by $n_\alpha = -\frac{\partial\Omega}{\partial\mu_\alpha}$. Then, using $\mu_a = \mu - h$ and $\mu_b = \mu + h$ in Eq. (43) we find

$$m = n_b - n_a = -\frac{d\Omega}{dh}, \tag{44}$$

and

$$n_T = n_b + n_a = -\frac{d\Omega}{d\mu}. \tag{45}$$

[1] S. Inouye *et al.*, Nature (London) **392**, 151 (1998).

[2] M. Greiner, C. A. Regal, and D. S. Jin, Nature (London) **426**, 537 (2003).

[3] S. Jochim *et al.*, Science **302**m 2101 (2003).

[4] M. W. Zwierlein *et al.*, Phys. Rev. Lett. **91**, 250401 (2003).

[5] G. Sarma, J. Phys. Chem. Solids **20**, 1029 (1963).

[6] W. V. Liu and F. Wilczek, Phys. Rev. Lett. **90**, 047002 (2003).

[7] P. Fulde and R. A. Ferrell, Phys. Rev. **135**, (1964); A. I. Larkin and Y. N. Ovchinnikov, Sov. Phys. JETP **20**, 762 (1965).

[8] P. F. Bedaque, H. Caldas, and G. Rupak, Phys. Rev. Lett. **91** 247002 (2003).

[9] H. Caldas, Phys. Rev. A **69** 063602 (2004).

[10] A. Sedrakian, J. Mur-Petit, A. Polls, and H. Muther, Phys. Rev. A **72** 013613 (2005).

[11] D. E. Sheehy and L. Radzihovsky, Phys. Rev. Lett. **96**, 060401 (2006); D. E. Sheehy and L. Radzihovsky, Ann. of Phys. **322**, 1790 (2007); L. Radzihovsky and D. E. Sheehy, Rep. Prog. Phys. **73**, 076501 (2010).

[12] M. W. Zwierlein, A. Schirotzek, C. H. Schunck, W. Ketterle, Science **311**, 492 (2006).

[13] M. W. Zwierlein, C. H. Schunck, A. Schirotzek, W. Ketterle, Nature (London) **442**, 54 (2006).

[14] G. B. Partridge, W. Li, R. I. Kamar, Y.-A. Liao, R. G. Hulet, Science **311**, 503 (2006).

[15] G. B. Partridge, W. Li, R. I. Karmar, Y. -A. Liao, R. G. Hulet, Phys. Rev. Lett. **97**, 190407 (2006).

[16] Y. Shin, C. H. Schunck, A. Schirotzek, W. Ketterle, Nature (London) **451**, 689 (2008).

[17] A. Gorlitz, J. M. Vogels, A. E. Leanhardt, C. Raman, T. L. Gustavson, J. R. Abo-Shaeer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle, Phys. Rev. Lett. **87**, 130402 (2001).

[18] V. Schweikhard, I. Coddington, P. Engels, V. P. Mogendorff, and E. A. Cornell, Phys. Rev. Lett. **92**, 040404 (2004).

[19] Y. Colombe, E. Knyazchyan, O. Morizot, B. Mercier, V. Lorent, H. Perrin, Europhys. Lett. **67**, 593 (2004).

[20] D. Rychtarik, B. Engeser, H.-C. Nagerl, and R. Grimm, Phys. Rev. Lett. **92**, 173003 (2004).

[21] J. Tempere, M. Wouters, and J. T. Devreese, Phys. Rev. B **75**, 184526 (2007).

[22] Lianyi He and Pengfei Zhuang, Phys. Rev. A **78**, 033613 (2008).

[23] G. J. Conduit, P. H. Conlon, and B. D. Simons, Phys. Rev. A **77**, 053617 (2008).

[24] T. N. De Silva, J. Phys. B **42**, 165301 (2009).

[25] X.-J. Liu, H. Hu, P. D. Drummond, Phys. Rev. A **76**, 043605 (2007).

[26] P. Kakashvili and C. J. Bolech, Phys. Rev. A **79**, 041603 (2009).

[27] Y. A. Liao, A. S. C. Rittner, T. Paprotta, W. Li, G. B. Partridge, R. G. Hulet, S. K. Baur, E. J. Mueller, Nature (London) **467**, 567 (2010).

[28] P. Pieri, D. Neilson, and G. C. Strinati, Phys. Rev. B **75**, 113301 (2007).

[29] K. Yamashita, K. Asano, and T. Ohashi, J. Phys. Soc. Jpn. **79**, 033001 (2010).

[30] M. A. Continentino, I. T. Padilha, J. Phys. Condens. Matter **20**, 095216 (2008); M. A. Continentino, I. T. Padilha, J. Phys. Condens. Matter **21**, 095603 (2009).

[31] M. Randeria, J.-M. Duan, and L.-Y. Shieh, Phys. Rev. Lett. **62**, 981 (1989).

[32] D. S. Petrov and G. V. Shlyapnikov, Phys. Rev. A **64**, 012706 (2001).

[33] V. M. Loktev *et al.*, Phys. Rep. **349**, 1 (2001), and references therein.

[34] H. Caldas, J. Stat. Mech. P11012 (2007).

[35] T. N. De Silva and E. J. Mueller, Phys. Rev. Lett. **97**, 070402 (2006).

[36] M. Haque and H. T. C. Stoof, Phys. Rev. Lett. **98**, 260406 (2006).

[37] S. K. Baur, S. Basu, T. N. De Silva, E. J. Mueller, Phys. Rev. A **79**, 063628 (2009).

[38] M. W. Zwierlein, private communication.

[39] A. Schirotzek, C.-H. Wu, A. Sommer, and M. W. Zwierlein, Phys. Rev. Lett. **102**, 230402 (2009).